EUROSOLIDS®

Humic Acids | Humic Acids Soluble

Euro Humic

Euro Humic is a highly concentrated, soluble source of Potassium Humate processed from leonardite via a unique extraction method. Our product is the most efficient source of Potassium Humate and has prominent abilities in regard to CEC, buffering and complexation. Euro Humic is recommended in poor soil conditions to enhance cation exchange capacity, and soil fertility. The presence of active humic extracts (long chain) in our formula has a positive effect on the uptake on all nutrients. Furthermore, it increases Nutrient Use Efficiency (NUE) and reduces leaching.



- Improves root and shoot growth with high quality buds
- Increases Cation-Exchange-Capacity of the soil
- Developed for fertigation in open field crops and for soil injection

Product Characteristics

- 85% of Humic Acids (ISO method)
- Wet-process method extracts long chain leonardite to improve efficiency of the Potassium Humate
- Patented micro granulation process
- Black granules
- Easy to handle, dissolve and apply

Compatibility

Compatible with other fertilizers, except Phosphate fertilizers. Therefore a separate tank is needed or fertilizers should be applied on different times.

Application

Composition (%w/w)

Potassium Oxide	12%
Total Humic Extracts	>85*| >75**%
Humic Acid	82*| 72**%
Fulvic Acid	> 3*| 3**%
Organic Nitrogen	1%
Ash content	31%
Dry matter	> 85%
Total Carbon	36%

Packaging

- 1kg
- 5kg
- 10kg

EUROSOLIDS®

Humic Acids

Euro Humic

Dosage | Fertigation

Crop	Application date	Min kg/ha/season	Max kg/ha/season
Vegetables	3 - 4 applications: • As of 1 week after planting • Until fruit setting • 10 - 14 days interval	8	16
Fruit trees (adult)	3 - 5 applications: • As of start of fertigation • Until fruit filling • 10 - 14 days interval	10	25
Citrus (adult)	4 - 6 applications: • As of start of fertigation • 10 - 14 days interval	15	25
Potato	3 - 5 applications: • As of start of fertigation • 10 - 14 days interval	8	20
Flowers	3 - 4 applications: • As of start of fertigation • 10 - 14 days interval	10	25
Vineyards (adult)	3 - 4 applications: • As of start of fertigation • 10 - 14 days interval	10	20
Banana	• During entire fertigation program • 10 - 14 days interval	10	30

The mentioned indicated dosages and application stages are given as a guideline. Exact dosages, concentration and application stage are subject to local conditions, use of other fertilizers and can only be given after an objective diagnosis.








Eurosolids Nederland | The Netherlands | tel.+31 (0)186 578 888






